Wildlife Best Management Practices (WBMPs) - Discussion Outline

- Make up of the Technical Advisory Committee (TAC) -Who drafted the WBMPs?
- Which state imperiled species "touch" the forest and forestry land uses?
- What do WBMPs look like?
- How does the WBMP process work?



Wildlife Best Management Practices for Forestry (WBMPs) - Technical Advisory Committee (TAC)

- The Technical Advisory Committee for forestry <u>water</u> <u>quality BMPs</u> is composed of: state and federal agencies, biologists, environmental groups, private & public landowners
- FWC did a <u>literature review on each species</u> and their habitat needs (part of the WBMP process)
- A <u>sub-group</u> of the TAC drafted the WBMPs; the <u>full TAC</u> committee reviewed, revised and approved the WBMPs



Wildlife Best Management Practices for Forestry (WBMPs) - What species are covered?

- Summary of state imperiled species in a forestry context:
 - Total number of species 16
 - Aquatic species 10 (62% of the total)
 - Bird species 4
 - Burrowing animal species 2
- Voluntary WBMPs are used by landowners that desire "coverage" under <u>incidental take</u> rules
- These <u>WBMPs</u> are for forestry (silviculture);
 not for development land uses



Wildlife Best Management Practices for Forestry (WBMPs) - How WBMPs work for <u>Aquatic Species</u>

- For 10 aquatic species (Crystal darter, Harlequin darter, Bluenose shiner, Blackmouth shiner, Tessellated darter, Santa Fe crayfish, Black Creek crayfish, Barbour's map turtle, Georgia blind salamander, Florida bog frog)
- Landowners that follow <u>water quality BMPs</u> are already <u>protecting aquatic habitat</u> by using Streamside Management Zones (SMZs)
- Landowners that <u>manage certified forests</u> (SFI, FSC, ATFS, etc.) <u>OR</u> send <u>wood to certified mills</u> are already required to follow water quality BMPs and protect T&E species



Wildlife Best Management Practices for Forestry (WBMPs) - How WBMPs work for <u>Bird Species</u>

- For 4 bird species (Little blue heron, Tricolored heron, Florida sandhill crane, Southeastern American kestrel aka sparrow hawk)
- Avoid heavy equipment operation within <u>330</u> of <u>heron</u>
 <u>rookeries</u> (Feb-May)
- Avoid heavy equipment operation within 400' of sandhill
 crane nests (found in non-forested wetlands) (Feb-May)
- Avoid heavy equipment operation within <u>490' of kestrel</u> nests (found in <u>open canopy uplands</u>) (Mar-Jun); retain snags



Wildlife Best Management Practices for Forestry (WBMPs) - How WBMPs work for <u>Burrowing Species</u>

- For 2 burrowing species (Gopher tortoise (GT) and Burrowing owl)
- Avoid heavy equipment concentrations near "known and visibly apparent" burrows;
- Burrows do <u>not</u> need to be flagged, mapped, etc.
- Avoid direct contact with burrows year round
- Avoid direct contact with burrow aprons May-Sep (GT eggs)
- Minimize heavy equipment during Sep-Oct (hatchling mortality)



Wildlife Best Management Practices for Forestry (WBMPs) - How WBMPs work generally

- All Wildlife BMPs (WBMPs) are voluntary
- These WBMPs only <u>apply to forestry uses</u> (silviculture) and <u>not development land uses</u>
- Landowners can enroll in WBMPs on a <u>species by species</u> basis and a tract by tract basis
- Landowners notify FDACS they intend to use WBMPs using a Notice of Intent (NOI) form (like water quality BMPs)
- Landowners are then presumed to be compliant with incidental take provisions when following WBMPs



Wildlife Best Management Practices for Forestry (WBMPs) - Timeline

- The subgroup TAC developed and revised the WBMPs in 2013/2014
- The <u>full TAC approved</u> the WBMPs in December, 2013
- Public hearing was held on June 20, 2014
- WBMPs may be available for landowner use soon
- Landowners that <u>manage certified forests</u> (SFI, FSC, ATFS, etc.) <u>OR</u> <u>send wood to certified mills</u> are already <u>required</u> to follow <u>water quality BMPs</u> and <u>protect T&E species</u>



Questions and Discussion





74# Wahoo

